

Effective Date: Monday, August 13, 2018

Test Updates

Immediate Action

In our continuing effort to provide you with the highest quality toxicology laboratory services available, we have compiled important changes regarding a number of tests we perform. Listed below are the types of changes that may be included in this notification, effective Monday, August 13, 2018

Test Changes - Tests that have had changes to the method/ CPT code, units of measurement, scope of analysis, reference comments, or specimen requirements.

Discontinued Tests - Tests being discontinued with alternate testing suggestions.

Please use this information to update your computer systems/records. These changes are important to ensure standardization of our mutual laboratory databases.

If you have any questions about the information contained in this notification, please call our Client Support Department at (866) 522-2206. Thank you for your continued support of NMS Labs and your assistance in implementing these changes.

The CPT Codes provided in this document are based on AMA guidelines and are for informational purposes only. NMS Labs does not assume responsibility for billing errors due to reliance on the CPT Codes listed in this document.



Test Updates

Test Code	Test Name	Test Name	Method / CPT Code		Stability	Scope	Reference Comments	Discontinue
2892B	Methimazole, Blood		•	•			•	
2892SP	Methimazole, Serum/Plasma		•	•			•	
2892U	Methimazole, Urine							•



Test Updates

Test Changes

loct enangee				
2892B Methimazole,	Methimazole, Blood			
Summary of Changes:	Specimen Requirements were changed. Specimen Requirements (Specimen Container) were changed. Reference Comment was changed. Methods/CPT Codes were changed [LC-MS/MS (80375)]			
Specimen Requirements:	1 mL Blood			
Transport Temperature:	Refrigerated			
Specimen Container:	Lavender top tube (EDTA)			
Light Protection:				
Special Handling:	•			
Rejection Criteria:	ria: Received Room Temperature.			
Scope of Analysis: Method (CPT Code)	ysis: LC-MS/MS (80375): Methimazole ode)			
Compound Name	Units	Reference Comment		
Methimazole	mcg/mL	Following single oral doses of 10 to 40 mg, average peak serum concentrations ranged from 0.2 to 0.8 mcg/mL at approximately 2 hours post dose.		
2892SP Methimazole,	Serum/Plasma			
Summary of Changes:	 Specimen Requirements were changed. Specimen Requirements (Specimen Container) were changed. Specimen Requirements (Special Handling) were changed. Reference Comment was changed. Methods/CPT Codes were changed [LC-MS/MS (80375)] 			

Sp	pecimen Requirements:	1 mL Serum or Plasma
	Transport Temperature:	Refrigerated
	Specimen Container:	Plastic container (preservative-free)
	Light Protection:	Not Required
	Special Handling:	Serum: Collect sample in Red top tube Plasma: Collect sample in Lavender top tube (EDTA) or Pink top tube. Promptly centrifuge and separate Serum or Plasma into a plastic screw capped vial using approved guidelines.
	Rejection Criteria:	Received Room Temperature. Polymer gel separation tube (SST or PST).
	Scope of Analysis: Method (CPT Code)	LC-MS/MS (80375): Methimazole
-		

Compound Name	Units	Reference Comment
Methimazole	mcg/mL	Following single oral doses of 10 to 40 mg, average peak serum concentrations ranged from 0.2 to 0.8 mcg/mL at approximately 2 hours post dose.



Test Updates

Discontinued Tests

Test Code	Test Name	Alternative Test
2892U	Methimazole, Urine	No Alternate Tests Available